

# Expanded Programme on Immunisation

Revised childhood immunisation schedule from December 2015

| Age of child                      | Vaccines needed  | How and where is it given |
|-----------------------------------|--|---------------------------|
| At birth                          | <b>BCG Bacilles Calmette Guerin</b>  | Right arm                 |
|                                   | <b>OPV (0) Oral polio vaccine</b>  | Drops by mouth            |
| 6 weeks                           | <b>OPV (1) Oral polio vaccine</b>  | Drops by mouth            |
|                                   | <b>RV (1) Rotavirus vaccine</b>  | Liquid by mouth           |
|                                   | <b>DTaP-IPV-Hib-HBV (1)</b><br>Diphtheria, tetanus, acellular pertussis, inactivated polio vaccine, haemophilus influenzae (type b) and hepatitis B combined | Intramuscular/Left thigh  |
|                                   | <b>PCV (1)</b><br>Pneumococcal conjugated vaccine  | Intramuscular/Right thigh |
| 10 weeks                          | <b>DTaP-IPV-Hib-HBV (2)</b><br>Diphtheria, tetanus, acellular pertussis, inactivated polio vaccine, haemophilus influenzae (type b) and hepatitis B combined | Intramuscular/Left thigh  |
| 14 weeks                          | <b>RV (2) Rotavirus vaccine *</b>  | Liquid by mouth           |
|                                   | <b>DTaP-IPV-Hib-HBV (3)</b><br>Diphtheria, tetanus, acellular pertussis, inactivated polio vaccine, haemophilus influenzae (type b) and hepatitis B Combined | Intramuscular/Left thigh  |
|                                   | <b>PCV (2)</b><br>Pneumococcal conjugated vaccine  | Intramuscular/Left thigh  |
| 6 months                          | <b>Measles vaccine (1) **</b>  | Subcutaneous/Left thigh   |
| 9 months                          | <b>PCV (3)</b><br>Pneumococcal conjugated vaccine  | Intramuscular/Right thigh |
| 12 months                         | <b>Measles vaccine (2) **</b>  | Subcutaneous/Right arm    |
| 18 months                         | <b>DTaP-IPV-Hib-HBV (4)</b><br>Diphtheria, tetanus, acellular pertussis, inactivated polio vaccine, haemophilus influenzae (type b) and hepatitis B combined | Intramuscular/Left arm    |
| 6 years<br>(Both boys and girls)  | <b>Td vaccine</b><br>Tetanus and reduced strength of diphtheria vaccine  | Intramuscular/Left arm    |
| 12 years<br>(both boys and girls) | <b>Td vaccine</b><br>Tetanus and reduced strength of diphtheria vaccine  | Intramuscular/Left arm    |

\*Rotavirus vaccine should NOT be administrated after 24 weeks.

\*\*Do not administer with any other vaccine.